

MULTI-TRANSONIC BLACK HOLE ACCRETION DISCS WITH ISOTHERMAL STANDING SHOCKS

TAPAS K. DAS^{1,2}, JAYANT K. PENDHARKAR^{3,4}, SANJIT MITRA³¹Division of Astronomy, Department of Physics and Astronomy, University of California at Los Angeles, Box 951562, Los Angeles, CA 90095-1562, USA² Institute of Geophysics and Planetary Physics, University of California at Los Angeles, Box 951567, Los Angeles, CA 90095-1567, USA³ IUCAA, Post Bag 4 Ganeshkhind, Pune 410 007, India⁴ Department of Astronomy, Osmania University, Hyderabad 500 007, India.
e-mail: tapas@astro.ucla.edu, jkp@iucaa.ernet.in, sanjit@iucaa.ernet.in

Draft version February 2, 2008

ABSTRACT

In this work we would like to address the issue of shock formation in black hole accretion discs. We provide a *generalized two parameter solution scheme* for multi-transonic, accretion and wind around Schwarzschild black holes, by mainly concentrating on accretion solutions which may contain steady, standing isothermal shocks. Such shocks conserve flow temperature by the dissipation of energy at the shock location. We use the vertically integrated 1.5 dimensional model to describe the disc structure, where the equations of motion apply to the equatorial plane of the central accretor, assuming the flow to be in hydrostatic equilibrium in the transverse direction. Unlike previous works in this field, our calculation is *not* restricted to any particular kind of post-Newtonian gravitational potentials, rather we use *all* available pseudo-Schwarzschild potentials to formulate and solve the equations governing the accretion and wind only in terms of flow temperature T and specific angular momentum λ of the flow. The accretion flow is assumed to be non-dissipative everywhere, except possibly at the shock location, if any. We observe that a significant region of parameter space spanned by $\{\lambda, T\}$ allows shock formation. Our generalized formalism assures that the shock formation is not just an artifact of a particular type of gravitational potential, rather inclusion of all available black-hole potentials demonstrates a substantially extended zone of parameter space allowing for the possibility of shock formation. We thus arrive at the conclusion that the standing shocks are essential ingredients in rotating, advective accretion flows of isothermal fluid around a non-spinning astrophysical black hole. We identify *all* possible shock solutions which may be present in isothermal disc accretion and thoroughly study the dependence of various shock parameters on fundamental dynamical variables governing the accretion flow for *all* possible initial boundary conditions. Types of shocks discussed in this paper may appear to be ‘bright’ because of the huge amount of energy dissipation at shock, and the quick removal of such energy to maintain the isothermality may power the strong X-ray flares recently observed to be emerged from our Galactic centre. The results are discussed in connection to other astrophysical phenomena of related interest, such as the QPO behaviour of galactic black hole candidates.

Subject headings: accretion, accretion disks — black hole physics — hydrodynamics — shock waves — QPO

1. INTRODUCTION

Perturbation of various kinds may produce discontinuities in astrophysical fluid flow. Discontinuities in fluid flow are said to take place over one or more surfaces when any dynamical and / or thermodynamic quantity may change discontinuously as such surfaces are crossed; the corresponding surfaces are called surfaces of discontinuity. Certain boundary conditions are to be satisfied across such surfaces and according to those conditions, surfaces of discontinuities are classified into various categories; the most important are being the shock waves or shocks, which, when the flow is adiabatic, obey the following conditions (Landau & Lifshitz, 1959, LL hereafter):

$$[\rho u] = 0, [p + \rho u^2] = 0, \left[\frac{u^2}{2} + w \right] = 0$$

where ρ, u, p and w are the fluid density, flow velocity, fluid pressure and the enthalpy of the flow respectively, and for any two arbitrary flow variables α and β ,

$$[\alpha + \beta] \equiv \{(\alpha + \beta)_- - (\alpha + \beta)_+\}$$

$$[\alpha\beta] \equiv \{\alpha_- \beta_- - \alpha_+ \beta_+\}.$$

The subscripts (+) and (−) refer to the quantities measured just after and just before the flow encounters the surface(s) of

discontinuity. Quite often, such shock waves are generated in various kinds of supersonic astrophysical flows having intrinsic angular momentum, resulting in a flow which becomes subsonic. This is because of the fact that the repulsive centrifugal potential barrier experienced by such flows are sufficiently strong to brake the motion and a stationary solution could be introduced only through a shock. Rotating, transonic astrophysical fluid flows are thus believed to be ‘prone’ to the shock formation phenomena. It has been established in recent years that in order to satisfy the inner boundary conditions imposed by the event horizon, accretion onto black holes should exhibit transonic properties in general; which further indicates that formation of shock waves are possible in astrophysical fluid flows onto galactic and extra-galactic black holes. One also expects that shock formation in black hole accretion discs might be a general phenomenon because shock waves in rotating astrophysical flows are convincingly able to provide an important and efficient mechanism for conversion of a significant amount of the gravitational energy (available from deep potential wells created by these massive compact accretors) into radiation by randomizing the directed infall motion of the accreting fluid. Hence shocks possibly play an important role in governing the

overall dynamical and radiative processes taking place in astrophysical fluids and plasma accreting onto black holes. Thus the study of steady, standing, stationary shock waves produced in black hole accretion has acquired a very important status in recent years.

The issue of the formation of steady, standing shock waves in black hole accretion discs is addressed in two different ways in general. First, one can study the formation of Rankine-Hugoniot shock waves (RHSW) in a polytropic flow. Radiative cooling in this type of shock is quite inefficient. No energy is dissipated at the shock and the total specific energy of the accreting material is a shock-conserved quantity. Entropy is generated at the shock and the post-shock flow possesses a higher entropy accretion rate than its pre-shock counterpart. The flow changes its temperature permanently at the shock. Higher post-shock temperature puffs up the post-shock flow and a quasi-spherical, quasi-toroidal centrifugal pressure supported boundary layer is formed in the inner region of the accretion disc. Fukue (1987) was the pioneer to initiate the systematic study of such shock dynamics using a rigorous equation of state valid for from the non-relativistic regime to the highly relativistic regime. Then Chakrabarti and collaborators (Chakrabarti 1989, 1996 and references therein, Abramowicz & Chakrabarti 1990, Chakrabarti and Molteni 1993) used the vertically integrated flow model of Matsumoto et. al. (1984) to extended Fukue's work towards the global study of shock formation in post-Newtonian geometry. However, such shock solutions were obtained either on a case by case basis, or, even when successful attempts were made to provide a more complete analysis, the boundary of the parameter space responsible for shock formation was obtained only for global variation of the total specific energy $\mathcal{E}$ (or accretion rate $\dot{M}$) and specific angular momentum λ of the flow, and not for variations of the polytropic constant γ of the flow; rather, accretion was usually considered to be ultra-relativistic¹, which may *not* always be a realistic assumption. As γ is expected to have great influence on the radiative properties of the flow in general, ignoring the explicit dependence of shock solutions on γ limits claims of generality. Also, all such so-called global shock solutions have been discussed only in the context of one particular type of pseudo-Schwarzschild black hole potential, namely, the Paczyński & Wiita (1980) potential (Φ_1 hereafter). For accretion in Kerr geometry, various authors (Lu 1985, 1986, Lu et al 1987, Nakayama and Fukue 1989) made attempts to provide a complete general relativistic description of shocked accretion flow by making a number of assumptions, some of which, however, may not appear to be fully convincing (see a detail discussion in Das 2002) and it is fair to say that no well accepted complete general relativistic global shock solution exclusively obtained for a hydrodynamic accretion disc around a Schwarzschild BH has yet appeared in the literature. Motivated by these limitations, Das (2002, D02 hereafter) provided a generalized formalism allowing one to construct and solve the equations governing the hydrodynamic black hole accretion of a polytropic fluid in all available pseudo-Schwarzschild potentials which may contain steady, standing RHSW. In this work, the explicit dependence of shock solutions on all important accretion parameters was thoroughly studied. The generalized formalism developed by D02 has also been used to explain related interesting phenomena like the

analytical computation of QPO frequencies (Das 2003) and the generation of accretion powered-galactic jets (Das, Rao and Vadawale 2003).

In another class of shock study, one concentrates on shock formation in isothermal black hole accretion discs. The characteristic features of such shocks are quite different from the non-dissipative shocks discussed above. In isothermal shocks, the accretion flow is allowed to dissipate a part of its energy and entropy at the shock surface to keep the post-shock temperature equal to its pre-shock value. This maintains the vertical thickness of the flow exactly the same just before and just after the shock is formed. Simultaneous jumps in energy and entropy join the pre-shock supersonic flow to its post-shock subsonic counterpart. Chakrabarti (1989a, C89 hereafter) studied the properties of such shocks in the rotating isothermal accretion around compact objects. However, his work was restricted to only one post-Newtonian black hole potential, i.e., the Paczyński and Wiita (1980) potential (Φ_1) and the dependence of shock parameters on initial boundary conditions was not thoroughly studied. Also, the degeneracy of the shock locations was not removed completely. Later on, Yang and Kafatos (1995, YK hereafter) investigated the formation of isothermal shocks and studied the stability properties of such shocks for general relativistic accretion onto Schwarzschild black holes. They also, quite successfully, computed the jump conditions for relativistic shocks. YK analyzed the stability of shocks in a more efficient way compared to the method used by C89 and could remove the two-fold degeneracy of shock locations; in this way they could properly identify the most stable shock location. However, unlike C89, YK used a one-dimensional conical model for accretion discs, where the existence of vertical hydrostatic equilibrium was not taken into account. Also, the explicit dependence of various shock-related quantities on accretion parameters was not fully explored. Nevertheless, YK still deserves special attention because it is the only work in the literature which attempts to study isothermal shock formation in a fully relativistic frame work.

In this paper, we would like to address the issue of shock formation in isothermal black hole accretion discs from a more general point of view. Instead of confining our calculation to a particular black hole potential, *all* available pseudo-Schwarzschild potentials will be exhaustively used to investigate the possibility of shock formation in hydrodynamic, inviscid, isothermal black hole accretion discs. Following Matsumoto et. al. (1984), we will be using the vertically integrated 1.5-dimensional model to describe the structure of the accretion discs, which is, perhaps, more realistic compared to simplified one dimensional conical flow model used by YK. In this 1.5 dimensional model (which has extensive use in literature, see Chakrabarti 1996 and references therein) the equations of motion are written on the equatorial plane of the central accretor whereas the flow remains in hydrostatic equilibrium in the transverse direction. For all black hole potentials we first formulate and solve the equations governing multi-transonic accretion and wind around the central accretor in terms of *only* two parameters, namely, specific flow angular momentum λ and flow temperature T . Then we provide a self-consistent scheme to capture the stable shock location(s) for multi-transonic accretion flow and finally we thoroughly

¹ By the term 'ultra-relativistic' and 'purely non-relativistic' we mean a flow with $\gamma = \frac{4}{3}$ and $\gamma = \frac{5}{3}$ respectively, according to the terminology used in Frank et. al. 1992.

study the dependence of various shock related quantities on fundamental accretion parameters. In this way we provide a generalized two parameter solution scheme for multi-transonic isothermal black hole accretion disc for all available pseudo-Schwarzschild black hole potentials which contains dissipative shock waves. Our generalized formalism assures that our model is not just an artifact of a particular type of potential only, and inclusion of every black hole potential allows a substantially extended zone of parameter space allowing for the possibility of shock formation.

Rigorous investigation of the complete general relativistic multi-transonic black hole accretion disc and wind is believed to be extremely complicated. At the same time it is understood that as relativistic effects play an important role in the regions close to the accreting black hole (where most of the gravitational potential energy is released), purely Newtonian gravitational potential (in the form $\Phi_N = -\frac{GM_{BH}}{r}$, where M_{BH} is the mass of the compact accretor) cannot be a realistic choice to describe transonic black hole accretion in general. To compromise between the ease of handling of a Newtonian description of gravity and the realistic situations described by complicated general relativistic calculations, a series of ‘modified’ Newtonian potentials have been introduced to describe the general relativistic effects that are most important for accretion disc structure around Schwarzschild black holes. Introduction of such potentials allows one to investigate the complicated physical processes taking place in disc accretion in a semi-Newtonian framework by avoiding pure general relativistic calculations so that most of the features of spacetime around a compact object are retained and some crucial properties of the analogous relativistic solutions of disc structure could be reproduced with high accuracy. Hence, those potentials might be designated as ‘pseudo-Schwarzschild’ potentials. Our calculations in this paper will be based on four such pseudo Schwarzschild potentials. Explicit forms of those four potentials are the following (see Das & Sarkar 2001, D02 and Artemova, Björnsson & Novikov 1996, ABN hereafter, and references therein, for detail discussion about various properties of these potentials):

$$\begin{aligned}\Phi_1(r) &= -\frac{1}{2(r-1)}; \quad \Phi_2(r) = -\frac{1}{2r} \left[1 - \frac{3}{2r} + 12 \left(\frac{1}{2r} \right)^2 \right] \\ \Phi_3(r) &= -1 + \left(1 - \frac{1}{r} \right)^{\frac{1}{2}}; \quad \Phi_4(r) = \frac{1}{2} \ln \left(1 - \frac{1}{r} \right)\end{aligned}\quad (1)$$

where r is the radial co-ordinate scaled in units of Schwarzschild radius. Hereafter, we will define the Schwarzschild radius r_g as

$$r_g = \frac{2GM_{BH}}{c^2}$$

(where M_{BH} is the mass of the black hole, G is universal gravitational constant and c is velocity of light in vacuum) so that the marginally bound circular orbit r_b and the last stable circular orbit r_s take the values $2r_g$ and $3r_g$ respectively for a typical Schwarzschild black hole. Also, total mechanical energy per unit mass on r_s (sometimes called ‘efficiency’ e) may be computed as -0.057 for this case. We will use a simplified geometric unit throughout this paper where radial distance r is scaled in units of r_g , radial dynamical velocity u and polytropic sound speed a of the flow is scaled in units of c (the velocity of light in vacuum), mass m is scaled in units of M_{BH} and all other derived quantities would be scaled accordingly. For simplicity, we will use $G = c = 1$. Among the above potentials, $\Phi_1(r)$ was

introduced by Paczyński and Wiita (1980) which accurately reproduces the positions of r_s and r_b . Also the Keplerian distribution of angular momentum obtained using this potential is exactly the same as that obtained in a pure Schwarzschild geometry. $\Phi_2(r)$ was proposed by Nowak and Wagoner (1991) to approximate some of the dominant relativistic effects of the accreting black hole (slowly rotating or non rotating) via a modified Newtonian potential. It has the correct form of r_s , and it produces the best approximation for the value of the angular velocity Ω_s (as measured at infinity) at r_s and the radial epicyclic frequency κ (for $r > r_s$). $\Phi_3(r)$ and $\Phi_4(r)$ were proposed by Artemova, Björnsson & Novikov 1996, ABN hereafter, to produce exactly the same value of the free-fall acceleration of a test particle at a given value of r as is obtained for a test particle at rest with respect to the Schwarzschild reference frame (Φ_3) and to produce the value of the free fall acceleration that is equal to the value of the covariant component of the three dimensional free-fall acceleration vector of a test particle that is at rest in the Schwarzschild reference frame (Φ_4) respectively. Hereafter, we will denote any i th potential as $\Phi_i(r)$ where $\{i = 1, 2, 3, 4\}$ corresponds to $\{\Phi_1(r), \Phi_2(r), \Phi_3(r), \Phi_4(r)\}$ respectively.

At this point, one may ask why we are interested in studying shock formation (and the overall structure of the isothermal black hole accretion disc in general) for various pseudo Schwarzschild potentials when YK have already performed shock study using complete general relativistic frame work. The answer is mainly two-fold. Firstly, we have used, perhaps, a more realistic flow geometry (1.5 dimensional vertically integrated disc model) than that of YK and we have provided a more detail dependence of various shock related quantities on fundamental accretion parameters. Secondly, and more seriously, even if someone can provide a completely satisfactory model for shock formation in full general relativistic isothermal black hole accretion, still the importance of our work may not be under estimated as we believe. Rigorous investigation of some of the shock related phenomena is extremely difficult (if not completely impossible) to study using full general relativistic framework. Hence one is expected to always rely on these pseudo-potentials because of the ease of handling them. Also, there are possibilities that in future someone may come up with a pseudo-Schwarzschild potential better than any of the potentials used in this paper, which will be the best approximation for complete general relativistic investigation of multi-transonic shocked flow. In such case, if one already formulates a generalized model for multi-transonic shocked isothermal accretion disc for any arbitrary black hole potential, exactly what we have done in this paper, then that generalized model will be able to readily accommodate that new $\Phi(r)$ without having any significant change in the fundamental structure of the formulation and solution scheme of the model and we need not have to worry about providing any new scheme exclusively valid only for that new potential, if any.

2. GOVERNING EQUATIONS

Hereafter we all use the notation $[\mathcal{P}_2]$, for a set of values of $\{\lambda, T\}$ for any particular Φ_i . It is to be mentioned here that for spherically symmetric isothermal accretion, one can solve the whole problem if only T alone is known. A one parameter solution scheme for Bondi (1952) type black hole accretion in all Φ_i s has also been formulated quite recently (Sarkar & Das

2002). For isothermal equation of state,

$$P = \frac{\rho RT}{\mu} = c_s^2 \rho \quad (2a)$$

where P, ρ, T and C_s are the pressure, density, constant temperature and the constant isothermal acoustic velocity of the flow respectively, R and μ being universal gas constant and the mean molecular weight, C_s and T can be related as

$$C_s = \Theta T^{\frac{1}{2}} \quad (2b)$$

where $\Theta = \sqrt{\frac{\kappa \mu}{m_H}}$ is a constant, $m_H \sim m_p$ being the mass of the hydrogen atom and κ is Boltzman's constant. Following standard literature, we consider a thin, rotating, axisymmetric, inviscid steady flow in hydrostatic equilibrium in transverse direction. The assumption of hydrostatic equilibrium is justified for a thin flow because for such flows, the infall time scale is expected to exceed the local sound crossing time scale in the direction transverse to the flow. The flow is also assumed to possess considerably large radial velocity which makes the flow 'advective'. We also assume that the disc to be non-self-gravitating, i.e. the increase of M_{BH} due to the accretion process itself is negligible. The local half-thickness $h_i(r)$ of the disc for any $\Phi_i(r)$ can be obtained by balancing the gravitational force by pressure gradient and can be expressed as

$$h_i(r) = \Theta \sqrt{\frac{rT}{\Phi'_i(r)}} \quad (3)$$

where $\Phi'_i(r) = \frac{d\Phi_i(r)}{dr}$. The radial momentum conservation equation and the equation of continuity could be written as:

$$u \frac{du}{dr} + \frac{1}{\rho} \frac{dP}{dr} + \frac{d}{dr} [\Phi_i^{eff}(r)] = 0 \quad (4a)$$

$$\frac{d}{dr} [u \rho r h_i(r)] = 0 \quad (4b)$$

where $\Phi_i^{eff}(r)$ is the effective potential which can be defined as the summation of the gravitational potential and the centrifugal potential for matter accreting under the influence of Φ_i and can be expressed as:

$$\Phi_i^{eff}(r) = \Phi_i(r) + \frac{\lambda^2(r)}{2r^2} \quad (5a)$$

However, for inviscid flow, $\lambda(r)$ has been assumed to be λ only. One can calculate the expression for the effective potential in full general relativity (D02):

$$\Phi_{GR}^{eff}(r) = r \sqrt{\frac{r-1}{r^3 - \lambda^2(1+r)}} - 1, \quad (5b)$$

to show that unlike $\Phi_i^{eff}(r)$, $\Phi_{GR}^{eff}(r)$ cannot be obtained by linearly combining the gravitational and rotational energy contributions because various energies in general relativity are combined together to produce non-linearly coupled new terms. Using eq. (2a - 2b) and (3), we integrate eq.(4a-4b) to obtain two conservation equations for our flow as:

$$\frac{u_e^2(r)}{2} + \Theta T \ln \rho_e(r) + \frac{\lambda^2}{2r^2} + \Phi_i(r) = C \quad (6a)$$

$$\dot{M}_{in} = \Theta \rho_e(r) u_e(r) r^{\frac{3}{2}} \sqrt{\frac{T}{\Phi'_i(r)}} \quad (6b)$$

Where C is a constant and $\dot{M}_{in}$ is the mass accretion rate. The subscript 'e' indicates the values measured on the equatorial

plane of the disc; however, we will drop 'e' hereafter if no confusion arises in doing so and will denote $\dot{M}_{in}$ by $\dot{M}$.

For a particular value of $[\mathcal{P}_2]_i$, it is quite straight forward to derive the space gradient of the dynamical flow velocity for any Φ_i as:

$$\frac{du}{dr} = \left[\frac{\left(\frac{3\Theta^2 T}{2r} + \frac{\lambda^2}{r^3} \right) - \left(\frac{1}{2} \Theta^2 T \frac{\Phi_i''(r)}{\Phi_i'(r)} + \Phi_i' \right)}{\left(u - \frac{\Theta^2 T}{u} \right)} \right] \quad (7)$$

where $\Phi_i'' = \frac{d^2 \Phi_i}{dr^2}$. Since the flow is assumed to be smooth everywhere, if the denominator of eq. (7) vanishes at any radial distance r , the numerator must also vanish there to maintain the continuity of the flow. One therefore arrives at the so called 'sonic point (alternately, the 'critical point') conditions' by simultaneously making the numerator and denominator of eq. (7) equal zero. The sonic point conditions can be expressed as:

$$u_s^i = \Theta T^{\frac{1}{2}} = \sqrt{\frac{\Phi_i'|_s - \frac{\lambda^2}{r_s^3}}{\frac{3}{2r_s} - \frac{1}{2} \left(\frac{\Phi_i''}{\Phi_i'} \right)_s}} \quad (8)$$

where the subscript 's' indicates the quantities are to be measured at the sonic point(s). For a fixed $[\mathcal{P}_2]_i$, one can solve the following polynomial of r_s to obtain the sonic point(s) of the flow:

$$\Phi_i''|_s + \frac{2}{\Theta^2 T} (\Phi_i')_s^2 - \left[\frac{3}{r_s \Theta} + \frac{2\lambda^2}{T \Theta^2 r_s^3} \right] \Phi_i'|_s = 0 \quad (9)$$

Solution of above equation is analogous to solving the following differential equation at critical points $x = x_c$:

$$\alpha \ddot{y} \Big|_{x=x_c} + [\{\beta \psi_1(x) + \gamma \psi_2(x)\}_{x=x_c} + \dot{y}|_{x=x_c}] \dot{y} \Big|_{x=x_c} = 0 \quad (10)$$

where :

$$y = f(x), \dot{y} = \frac{dy(x)}{dx}, \ddot{y} = \frac{d^2 y(x)}{dx^2},$$

$f(x), \psi_1(x)$ and $\psi_2(x)$ are different functions of x and $\{\alpha, \beta, \gamma\}$ are either constant or various functions of initial parameters.

The critical value of the dynamical velocity gradient at sonic point(s) can be computed by solving the following equation for $\left(\frac{du}{dr}\right)_{s,i}$:

$$\left(\frac{du}{dr}\right)_{s,i} = \pm \frac{1}{\sqrt{2}} \left[\frac{1}{2} \Theta^2 T \left[\left(\frac{\Phi_i''}{\Phi_i'} \right)_s^2 - \left(\frac{\Phi_i'''}{\Phi_i'} \right)_s \right] - \left(\Phi_i''|_s + \frac{3\Theta^2 T}{2r_s^2} + \frac{3\lambda^2}{r_s^4} \right) \right]^{\frac{1}{2}} \quad (11)$$

3. MULTI-TRANSONIC ACCRETION AND WIND

For all Φ_i , we find a significant region of parameter space spanned by $[\mathcal{P}_2]_i$ which allows the multiplicity of sonic points for accretion as well as for wind where two real physical inner and outer (with respect to the black hole location) X type sonic points r_{in} and r_{out} encompass one 'O' type unphysical middle sonic point r_{mid} in between. For a fixed set of $\{\lambda, T\}$, one can solve eq. (9) to obtain required sonic points and for a particular set of values of $\{\lambda, T\}$, one obtains at most four real roots, one of which is always less than one r_g , hence falls inside the event horizon. Thus maximum number of real physical roots

is three, hence maximum number of sonic points for any $[\mathcal{P}_2]_i$ is also three. Apparently there remains an ambiguity in determining the values of $[\mathcal{P}_2]_i$ for which three sonic points form in accretion and the values of $[\mathcal{P}_2]_i$ for which three sonic points form in wind because just by studying the values of $\{\lambda, T\}$ or $\{r_{in}, r_{mid}, r_{out}\}$, one cannot determine whether it is accretion or wind which is multi-transonic for that $[\mathcal{P}_2]_i$. However, we can unambiguously classify the parameter space in the following way:

Let us suppose that for a particular value of $[\mathcal{P}_2]_i$, accretion can be multi-transonic. So the subsonic far field solution will become supersonic after crossing the outer sonic point r_{out}^{acc} in accretion. It would manifest the tendency for passing through the inner sonic point r_{in}^{acc} again after a shock transition if the solution passing through the inner sonic point is more preferred by nature for some physical reason(s). One can show that it is the total energy content of the flow which will discriminate the solutions passing through r_{out}^{acc} and r_{in}^{acc} . If the energy content of the solution passing through r_{in}^{acc} is lower than the solution passing through r_{out}^{acc} , then accretion flow will prefer to pass through r_{in}^{acc} because nature will always prefer a lower energy solution. The whole situation would be reversed for multi-transonic winds, energy content of wind solutions through r_{in}^{wind} will be *higher* than that through r_{out}^{wind} . One can calculate the energy difference between solutions passing through r_{out} and r_{in} as:

$$\delta\epsilon = \epsilon_{out} - \epsilon_{in} = \frac{\lambda^2}{2} \left(\frac{1}{r_o^2} - \frac{1}{r_i^2} \right) + (\Phi_i|_{r_o} - \Phi_i|_{r_i}) + \ln \left[\left(\frac{r_i}{r_o} \right)^{\frac{3}{2}} \left(\frac{\Phi_i'|_{r_o}}{\Phi_i'|_{r_i}} \right)^{\frac{1}{2}} \right]^{\Theta^2 T} \quad (12)$$

where $r_o = r_{out}$ and $r_i = r_{in}$. For a particular Φ_i , if $\mathcal{A}_i[\mathcal{P}_2]_i$ denotes the universal set representing the entire parameter space covering all values of $[\mathcal{P}_2]_i$, and if $\mathcal{B}_i[\mathcal{P}_2]_i$ represents one particular subset of $\mathcal{A}_i[\mathcal{P}_2]_i$ which contains only the particular values of $[\mathcal{P}_2]_i$ for which eq. (9) gives three real physical roots (four real roots with one unphysical root lying inside the event horizon), then $\mathcal{B}_i[\mathcal{P}_2]_i$ can further be decomposed into two subsets $\mathcal{C}_i[\mathcal{P}_2]_i$ and $\mathcal{D}_i[\mathcal{P}_2]_i$ such that

$$\mathcal{C}_i[\mathcal{P}_2]_i \subseteq \mathcal{B}_i[\mathcal{P}_2]_i \text{ only for } \delta\epsilon > 0$$

and:

$$\mathcal{C}_i[\mathcal{P}_2]_i \subseteq \mathcal{D}_i[\mathcal{P}_2]_i \text{ only for } \delta\epsilon < 0$$

For $[\mathcal{P}_2]_i \in \mathcal{C}_i[\mathcal{P}_2]_i$, we get multi-transonic *accretion* and for $[\mathcal{P}_2]_i \in \mathcal{D}_i[\mathcal{P}_2]_i$ one obtains multi-transonic *wind*. In Fig. 1, we plot:

$$(\lambda_i, T_i) \in [\mathcal{P}_2]_i \in \mathcal{C}_i[\mathcal{P}_2]_i \subseteq \mathcal{B}_i[\mathcal{P}_2]_i$$

and in Fig. 2. we plot:

$$(\lambda_i, T_i) \in [\mathcal{P}_2]_i \in \mathcal{D}_i[\mathcal{P}_2]_i \subseteq \mathcal{B}_i[\mathcal{P}_2]_i$$

For all Φ_i , while the specific angular momentum is plotted along X axis, the flow temperature T_{10} , scaled in the unit of $10^{10}K$, is plotted along Y axis. For Fig. 1, $\mathcal{A}_i\mathcal{B}_i\mathcal{C}_i$ represents the region $\mathcal{C}_i[\mathcal{P}_2]_i$ for a particular Φ_i and for Fig. 2, $\mathcal{A}_i\mathcal{B}_i\mathcal{C}_i\mathcal{D}_i$ represents the region $\mathcal{D}_i[\mathcal{P}_2]_i$ for a particular Φ_i . From Fig. 1 it is evident that no region of parameter space common to all Φ_i is found for which $[\mathcal{P}_2]_i \in \mathcal{C}_i[\mathcal{P}_2]_i$. However, significant region of parameter space is obtained for which $[\mathcal{P}_2]_i \in \mathcal{C}_i[\mathcal{P}_2]_i$ for Φ_2 and Φ_3 . It is interesting to note that (see Fig. 2 of D02) same trend is observed for multi-transonic *polytropic* accretion as well. The maximum flow temperature in T_{10} for which one

can obtain multi-transonic accretion may be denoted by T_i^{max} for any $\Phi_i(r)$. One observes that

$$T_3^{max} > T_4^{max} > T_1^{max} > T_2^{max}$$

and the minimum value of angular momentum λ_i^{min} for which one obtains the multi-transonic accretion, can be arranged as:

$$\lambda_3^{min} < \lambda_2^{min} < \lambda_4^{min} < \lambda_1^{min}$$

Hence one can observe that for Φ_3 and Φ_2 , even very low angular momentum flow (Quasi-Bondi accretion, so to say) may produce multi-transonic accretion.

Situation is not exactly similar for multi-transonic *wind*. From Fig. 2 one observes that there is a common region for $\mathcal{D}_i[\mathcal{P}_2]_i$ for flows in Φ_1 , Φ_4 and Φ_3 , and there exist a separate common zone for $\mathcal{D}_i[\mathcal{P}_2]_i$ in between Φ_2 , Φ_4 and Φ_3 . Here also same sequential order is maintained for T_i^{max} :

$$T_3^{max} > T_4^{max} > T_1^{max} > T_2^{max}$$

and for λ_i^{min} :

$$\lambda_3^{min} < \lambda_2^{min} < \lambda_4^{min} < \lambda_1^{min}$$

However, there are significant difference for range of λ for multi-transonic accretion and wind. Unlike multi-transonic accretion, three sonic points in *wind* form for considerably high values of λ_i ; particularly for Φ_1 and Φ_4 , the value of λ is very large. It is to be noted that for *polytropic* accretion and wind, the situation was quite different (see Fig. 2 of D02); unlike isothermal accretion and wind, it was observed there that

$$\lambda_i^{max} \Big|_{\text{accretion}} \sim \lambda_i^{max} \Big|_{\text{wind}}$$

It is evident from Fig. 2 that multi-transonic wind is usually obtained only for strongly rotating flows, i.e., for flows with intense rotational energy content, and quasi-Bondi flow does not produce multi-transonic wind, this trend is more explicit for flows in Φ_1 and Φ_4 .

4. SHOCK FORMATION AND RELATED PHENOMENA

In this work we are not interested in shock formation in wind, rather we will concentrate on shock formation in accretion only. For any Φ_i , if shock forms, then $[\mathcal{P}_2]_i$ responsible for shock formation must be somewhere from the region $\mathcal{A}_i\mathcal{B}_i\mathcal{C}_i$ (see Fig. 1), i.e, for which $[\mathcal{P}_2]_i \in \mathcal{C}_i[\mathcal{P}_2]_i$, though not all such $[\mathcal{P}_2]_i$ will allow shock transition. A multi-transonic accretion solution containing shock waves may behave as follows:

The far-field subsonic solution will become supersonic after passing the outer sonic point r_{out} . As the specific energy content of the solution passing through the inner sonic point r_{in} is less compared to the already supersonic solution through r_{out} , the solution through r_{out} will tend to 'jump down' to its counter solution passing through r_{in} . This jump will take place via a shock transition and certain amount of energy will be dissipated at shock, while the flow temperature (hence the acoustic flow velocity) will remain the same. The post shock flow will become subsonic and will again accelerate to supersonic as it passes through r_{in} . The combined shocked flow field is thus subsonic at infinity and supersonic near the black hole, a physically acceptable solution for black hole accretion.

Like temperature, the baryon flux, the momentum flux and a specific function of Mach number are also conserved at shock. Following C89, those conservation equation could be written as:

$$\left[\rho M \Theta T^{\frac{1}{2}} \right] = 0, [P + \rho u^2] = 0, \left[M + \frac{1}{M} \right] = 0 \quad (13)$$

where M denotes the Mach number, and ρ and P are the vertically integrated values of density and pressure. Also

$$[\alpha + \beta] \equiv \{(\alpha + \beta)_- - (\alpha + \beta)_+\}$$

$$[\alpha\beta] \equiv \{\alpha_- \beta_- - \alpha_+ \beta_+\}$$

as described in §1. Using eq. (13), we can formulate a generalized shock condition, which is the following

$$\frac{1}{\rho_-} - \frac{u_-^2 + \Theta^2 T}{u_-^2 \rho_- + (2\rho_+ - \rho_-)\Theta^2 T} = 0 \quad (14)$$

The above equation is valid *only* at the shock location, "+" and "-" refers to the post and pre shock quantities respectively. For the kind of shock discussed here, the shock strength S_i (which is basically the ratio of the pre- to post-shock flow Mach number) is exactly equal to the shock compression ratio R_{comp} (the ratio of vertically integrated post- to pre-shock flow density).

$$S_i = R_{comp} \quad (15)$$

The amount of energy dissipation at shock can be computed as

$$\Delta\epsilon = \frac{\dot{M}^2 \Phi_i'}{\Theta^4 T^2 \rho_+^2 r_{sh}^3} \Big|_{r=r_{sh}} (R_{comp}^2 - 1) - \Theta^2 T \ln(R_{comp}) \quad (16)$$

where $\dot{M}$ is the accretion rate scaled in the unit of Eddington rate. In another way eq. (16) can be written in $\dot{M}$ independent form as:

$$\Delta\epsilon = \frac{1}{2} (M_-^2 - M_+^2) - \Theta^2 T \ln S_i \quad (17)$$

Fig. 3 demonstrates few typical flow topologies of the integral curves of motion for shocked flow in various Φ_i (indicated in the figure). While the distance from the event horizon of the central black hole (scaled in the units of r_g and plotted in logarithmic unit) is plotted along the X axis, the local Mach number of the flow is plotted along the Y axis. One can easily obtain such a set of figures for any $[\mathcal{P}_2]_i$ which allow shock transition. For all figures, ABC_1C_2D represents the transonic accretion passing through the outer sonic point r_{out} (marked as B) if a shock would not form. However, as $\delta\epsilon$ in between r_{out} and r_{in} is greater than zero, the flow will encounter a shock, becomes subsonic and jumps on the branch JEF, which ultimately hits the event horizon supersonically after it passes through the inner sonic point r_{in} marked by $\mathcal{I}$ in the figure. The small circle with a dot at its centre ("⊙") indicates the location of the unphysical middle sonic point r_{mid} . It is to be mentioned that by solving eq. (14) one obtains *two* formal shock locations C_1 and C_2 (vertical lines marked by arrowheads through C_1 and C_2 represent the shock transitions). For every Φ_i and for any $[\mathcal{P}_2]_i$ allowing shock transition, r_{mid} *always* lies in between two shocks passing through C_1 and C_2 . In other words, the 'outer' shock at C_1 is obtained in between r_{out} and r_{mid} while the 'inner' shock at C_2 is obtained in r_{mid} and r_{in} . For all Φ_i s, it is observed that the location of the inner and the outer sonic points anti-correlate with λ as well as with T . Hotter and strongly rotating flows push the real physical sonic points towards the event horizon. The corresponding values of $\{r_{in}, r_{mid}, r_{out}\}$, the shock locations at C_1 and C_2 and the corresponding shock strengths are indicated at the top of each figure, while the corresponding values of λ and T (scaled in the unit $T_{10} \rightarrow T/T_{10} (= 10^{10} \text{K})$) for which the solutions are obtained, are indicated inside each figure. GBH represents the 'self wind' of the flow, which, in the course of it's motion away from the black hole to infinity, becomes supersonic after passing through r_{out} at B. The overall

scheme for obtaining the above mentioned integral curves is as follows:

For a particular $[\mathcal{P}_2]_i$, we first compute r_{in} , r_{mid} and r_{out} by solving eq. (9). Then we obtain the dynamical velocity gradient of the flow at sonic points by solving eq. (11). We then calculate the local dynamical flow velocity $u(r)$, the local radial Mach number $M(r)$, the fluid density $\rho(r)$ and any other related dynamical or thermodynamic quantities by solving the eq. (6 - 11) from the outer sonic point using fourth order Runge-Kutta method. We start integrating from r_{out} in two different directions. Along BH, we only solve for $u(r)$ and $M(r)$, because shock does not form in subsonic flows. Integration along BC_1C_2D involves a different procedure. Along BC_1C_2D , along with our calculation of $u(r)$ and $M(r)$, we also, at every integration step (with smallest possible step size), keep checking whether eq. (14) is being satisfied at that point. Along the curve $\mathcal{IJ}$ we calculate ρ_+ and u_+ . At the same time, along the curve BC_1C_2D , from the initial point whose r coordinate is equal to the r coordinate of J , to the final point whose r coordinate is equal to r_{in} , we calculate ρ_- . Then we substitute these values of ρ_- , ρ_+ and u_- in eq. (14) and check for which r coordinate the values of ρ_- , ρ_+ and u_- satisfy the equation. We then pick up that particular point as our shock location r_{sh} and calculate the value of S_i and R_{comp} for the shock location(s).

Like C89 and YK, we also obtain two real physical formal shock locations C_1 and C_2 . C89 obtained exactly same shock strength for all formal shock locations. It is important to note that their conclusion is *not* true. For *all* Φ_i and for all values of $[\mathcal{P}_2]_i$ allowing shock formation, we obtain *distinctively different* shock strength for shocks through C_1 and C_2 . We did not follow the stability analysis provided by C89 because that kind of stability analysis cannot remove the degeneracy of shock locations, it can only partially remove the multiplicity. We follow a more robust stability analysis provided by YK where a *unique* stable shock location was obtained. Following YK, we find that only the shock through C_1 , i.e. the 'outer' shock which forms in between r_{out} and r_{mid} is stable, while the other one (shock through C_2) is not. It is important to note that Nakayama (1992) also made important contribution to remove the degeneracy of multiple shocks in relativistic flow. Hereafter, we will deal with only the stable 'outer' shock location. In Fig. 4 we represents the *entire* parameter space spanned by $[\mathcal{P}_2]_i$ which allows shock formation for *all* Φ_i . The specific flow angular momentum λ is plotted along the X axis while the flow temperature T (scaled in the unit of $T_{10} \rightarrow 10^{10} \text{K}$) is plotted along the Y axis. For any pseudo-potential Φ_i , $A_iB_iC_i$ represents the parameter space for which $[\mathcal{P}_2]_i \in C_i[\mathcal{P}_2]_i$. The region bounded by $D_iB_iC_i$ and marked by S_i represents the parameter space for which the stable standing shock forms for that particular Φ_i and the region $A_iB_iD_i$ represents the region for which accretion is multi-transonic yet no stable standing shock forms in accretion. If $[\mathcal{P}_2]_i \in \mathcal{E}_i[\mathcal{P}_2]_i \subseteq C_i[\mathcal{P}_2]_i$ represents the parameter space for stable standing shock formation, obviously:

$$D_iB_iC_i \equiv [\mathcal{P}_2]_i \in \mathcal{E}_i[\mathcal{P}_2]_i \subseteq C_i[\mathcal{P}_2]_i \quad (18)$$

One can also define $[\mathcal{P}_2]_i \in \mathcal{F}_i[\mathcal{P}_2]_i$ which is complement of $\mathcal{E}_i[\mathcal{P}_2]_i$ related to $C_i[\mathcal{P}_2]_i$, so that:

$$A_iB_iD_i \equiv \left[\mathcal{F}_i[\mathcal{P}_2]_i \mid [\mathcal{P}_2]_i \in C_i[\mathcal{P}_2]_i, [\mathcal{P}_2]_i \notin \mathcal{E}_i[\mathcal{P}_2]_i \right] \quad (19)$$

The shock location becomes imaginary in $\mathcal{F}_i[\mathcal{P}_2]_i$ hence no stable shock forms in the region $A_iB_iD_i$ even if for $[\mathcal{P}_2]_i \in A_iB_iD_i$

produces multi-transonic accretion. Hence one should remember that although the multiplicity of sonic points in accretion is a necessary condition for the existence of shocks, but is *not* a sufficient one.

We calculate $r_{sh}, \Delta\epsilon, S_i, R_{comp}$ and all possible shock related quantities for $[\mathcal{P}_2]_i \in D_i B_i C_i$ for all Φ_i s. Hence our calculation is much more general one compared to C89 or YK. In Fig. 5, we show the variation of r_{sh} (plotted along the Z axis) with λ (plotted along the X axis) and T (scaled in the unit of T_{10} and plotted along the Y axis) for all four Φ_i ($\Phi_1 \rightarrow (a), \Phi_2 \rightarrow (b), \Phi_3 \rightarrow (c), \Phi_4 \rightarrow (d)$).

One observes that the maximum possible value of shock location is obtained for flows in Φ_1 while Φ_3 allows minimum span in the value of r_{sh} . In Fig. (6), we plot the variation of the energy dissipation at shock, i.e. the value of $\Delta\epsilon$ (plotted along Z axis) with the flow angular momentum λ (plotted along the X axis) and the flow temperature T (scaled in the unit of T_{10} and plotted along the Y axis) for all four pseudo potentials marked in the figure as $\Phi_1 \rightarrow (A), \Phi_2 \rightarrow (B), \Phi_3 \rightarrow (C), \Phi_4 \rightarrow (D)$. The shock location correlates with λ as well as with T . This is obvious because a rapidly rotating flow will generate a centrifugal barrier at a relatively large distance from the event horizon and a hotter flow will generate sufficient amount of thermal pressure to break the flow well away from the black hole. Also it is observed that the stronger the shock (shock with large compression ratio), the higher the amount of dissipated energy at the shock location and that a shock formed closer to the event horizon will result in a larger amount of energy dissipation. This is also obvious because the closer the shock location to the event horizon, the higher the amount of available gravitational potential energy to be dissipated at the shock.

5. CONCLUDING REMARKS

The existing literature on shock formation in black hole accretion discs are heavily inclined towards the study of polytropic accretion and non-dissipative shocks while the formation of dissipative shocks in an isothermal flow is a relatively less well studied subject. Such a ‘preferential investigation’ is perhaps due to the fact that global isothermality in ‘black hole accretion’ is difficult to achieve for realistic flows (it is difficult to maintain exactly the same temperature throughout a disc), the more realistic kind of accretion is expected to be described using a polytropic equation of state. However, one should never forget that the techniques developed to study disc geometry and shock dynamics in all existing literature (including the present paper) are heavily based on various assumptions, and hence, are far from being sufficiently complete and self-consistent to model the exact physical situation. Even if the vertically integrated 1.5 dimensional disc model developed by Matsumoto et. al. (1984), which has been used in this paper, is believed to be one of the most successful disc models, it suffers from a number of limitations in studying the polytropic shock dynamics. One of such major limitations is, even for an extremely thin shock, i.e. a polytropic shock with effective zero thickness, the thickness of the flow (i.e. the disc thickness) changes abruptly at the shock (pre and post shock half thickness of the polytropic flow is quite different) and it is impossible to write down the exact pressure balance condition at the shock by vertical averaging. On the other hand, for isothermal flows with dissipative shocks (the kind of shock dynamics studied in this paper) at constant acoustic velocity, the flow thickness remains the same (because of the quick removal of dissipated energy)

and a correct pressure balance condition is obtained. Hence we believe that the study of shock formation in isothermal flow is quite important in its own right and we have provided *for the first time* a generalized formalism to study such shocks for *all* available pseudo-Schwarzschild black hole potentials. Also we have studied the *complete* disc dynamics for multi-transonic accretion and wind and have demonstrated *all* the shock solutions and successfully studied the dependence of important shock quantities on every fundamental accretion parameter. However, in this work it has not been possible to compare the different potentials in connection to the suitability of mimicking the exact general relativistic solutions as was done in D02. It was beyond the scope of this paper to formulate and solve the complete general relativistic set of equations governing the isothermal accretion and wind, we hope to present that work somewhere else.

It is to be mentioned here that a number of recent works study the dynamical and radiative properties of the post-shock region of advective accretion discs (inner part of the disc, so to say) in order to understand the spectral properties of the black hole candidate (Shrader & Titarchuk 1998, and references therein) and to theoretically explain a number of diverse phenomena which are supposed to take place in the inner part of the shocked black hole accretion discs. (Titarchuk, Lapidus & Muslimov 1998, and references therein, Das 1998, Das & Chakrabarti, 1999, Das 2003, Das, Rao & Vadawale, 2003, Naik et.al. 2001). The importance of studying the physics of post-shock accretion flow has also been realized from observational evidence (Rutledge et. al. 1999, Muno, Morgan & Remillard 1999, Webb & Malkan 2000, Smith, Heindl and Swank 2002). Thus we believe that our generalized model presented in this paper may have far reaching consequences in handling various shock related phenomena because this model deals with all available black hole potentials. Nevertheless, one limitation of our model is that we have assumed the flow to be totally inviscid and viscous dissipation of azimuthal angular momentum was not taken into account. Exact modeling of viscous multi-transonic black hole accretion, including proper heating and cooling mechanisms, is quite an arduous task, and we did not dare to attempt that in this paper. However, qualitative calculations show that the introduction of viscosity via a radius-dependent power-law distribution for angular momentum only weakens the strength of the centrifugal barrier and pushes the shock location closer to the event horizon, keeping the overall basic physics concerning the shock dynamics and related issues unaltered; details of this work is in progress and will be discussed elsewhere.

In our work, some of the examples of the shocked multi-transonic flow, especially for flows in Φ_2 and Φ_3 , contain low intrinsic angular momentum. Such weakly rotating flows may be found in nature in systems like detached or semi-detached binaries fed by accretion from OB stellar winds (Illarionov & Sunyaev 1975, Liang & Nolan 1984), semi-detached low-mass non-magnetic binaries (Bisikalo et al. 1998) or super-massive black holes fed by accretion from slowly rotating central stellar clusters (Illarionov 1988, Ho 1999 and references therein).

It is now a well established fact that a number of galactic black hole candidate show QPO behaviours. Associated large amplitudes and short time scales of such intensity variation

suggests that such oscillations are a diagnostic of the accretion processes in the innermost regions of the accretion discs of the central accretor (van der Kluis, 1989, and references therein) and it has been suggested in recent years that such QPOs may be generated because of the shock oscillation in black hole accretion discs (Molteni, Sponholz & Chakrabarti 1996, Titarchuk, Lapidus & Muslimov 1999). We apply our shock solutions obtained in this paper to compute the frequency of the QPOs generated by such shock oscillations (Das 2003) and study the dependence of such frequencies on various fundamental accretion and shock parameters.

We would like to complete our discussion by pointing out one important finding in our work. Since the accretion flows discussed in this paper are chosen to be inviscid and non-dissipative, most of the binding energy of the flow is dissipated at the shocks. Thus such shocks can serve as the efficient provider of effective radiative cooling mechanism in the accretion discs. From Fig. 6 it is evident that for some region of parameter space spanned by $[\mathcal{P}_2]_i$, the amount of energy dissipated at the shock could be as high as more than ten percent of the total rest mass energy of the flow. This large amount of energy has to be immediately removed from the accretion disc (at least from the equatorial plane) to maintain the isothermality of the flow. This dissipated energy may be drained from the disc through various processes like a sudden X-ray burst (flare) of energy from the accretion discs close to the event horizon (few r_g away) or through energetic leptonic/ baryonic outflows

and jets, the formation of which is likely to be initiated by this huge amount of dissipated energy at the shock. Observational signature of such flares from the close vicinity of our Galactic centre black hole has already been reported quite recently (Baganoff et al. 2001). The observed length scale of the accretion flow from where such flares are generated, are in fairly close agreements with our theoretical calculations presented in this paper. Thus one concludes that the kind of shock discussed in this paper could be efficiently radiative and ‘bright’. The energy dissipation at the shock and its channeling from the accretion disc system may explain some interesting related issues like shock generated X-ray flares and the QPO of galactic microquasars (Das & Rao 2003, in preparation) and the generation mechanism of accretion powered galactic jet (Das 2003a, in preparation).

This research has made use of NASA’s Astrophysics Data System Bibliographic Services. Research of TKD at UCLA is supported by NSF funded post doctoral fellowship (Grant No. NSF AST-0098670). SM acknowledges the financial assistance from CSIR (India) in the form of graduate student fellowship. JKP would like to acknowledge the kind hospitality provided by IUCAA through the visiting student research program. Authors owe much debt to Prof. Mark R. Morris for reading the manuscript extremely carefully and for providing a number of useful suggestions. Authors would also like to thank the anonymous referee and the editor Prof. Luigi Stella for useful comments, and Atashi Chatterjee for typing the manuscript.

REFERENCES

- Abramowicz, M., & Chakrabarti, S. K. 1990, *ApJ*, 350, 281
 Artemova, I. V., Björnsson, G., & Novikov, I. D. 1996, *ApJ*, 461, 565 (ABN)
 Baganoff, F. et al. 2001, *Nature*, Vol 413, Pg. 45
 Bisikalo, A. A., Boyarchuk, V. M., Chechetkin, V. M., Kuznetsov, O. A., & Molteni, D. 1998, *MNRAS*, 300, 39
 Bondi, H. 1952, *MNRAS*, 112, 195
 Chakrabarti, S. K. 1989, *ApJ*, 347, 365
 Chakrabarti, S. K. 1989a, *MNRAS*, 240, 7 (C89)
 Chakrabarti, S. K. 1996, *Physics Reports*, 266, 229
 Chakrabarti, S. K., & Molteni, D. 1993, *ApJ*, 417, 671
 Das, T. K. 1998, in *Observational Evidence for Black Holes in the Universe*, Ed. S. K. Chakrabarti (Kluwer Academic: Holland), p. 113
 Das, T. K., 2002, *ApJ*, 577, 880
 Das, T. K., 2003, *ApJL*, To Appear (astro-ph/0302014)
 Das, T. K., 2003a, In Preparation.
 Das, T. K., & Chakrabarti, S. K. 1999, *Class. Quantum Grav*, 16, 3879
 Das, T. K., & Rao, A. R., 2003, In Preparation.
 Das, T. K., & Sarkar, A. 2001, *A & A*, 374, 1150
 Das, T. K., Rao, A. R., & Vadawale, S. 2003, *MNRAS*, To Appear (astro-ph/0301344)
 Frank, J., King, A., & Raine, D. 1992, *Accretion Power in Astrophysics*. 2nd. Edition. Cambridge University Press.
 Fukue, J. 1987, *PASJ*, 39, 309
 Ho, L. C. 1999, in *Observational Evidence For Black Holes in the Universe*, ed. S. K. Chakrabarti (Dordrecht: Kluwer), 153
 Illarionov, A. F. 1988, *Soviet Astron.*, 31, 618
 Illarionov, A.F., & Sunyaev, R. A. 1975a, *A & A*, 39, 205
 Landau, L. D., & Lifshitz, E. D. 1959, *Fluid Mechanics*, New York: Pergamon (LL)
 Liang, E. P. T., & Nolan, P. L. 1984, *Space Sci. Rev.*, 38, 353
 Lu, J. F. 1985, *A & A*, 148, L176
 Lu, J. F. 1986, *GReGr*, 18, L45
 Lu, J. F., Yu, K. N., Yuan, F., & Young, E. C. M. 1997, *A & A*, 321, 665
 Matsumoto, R., Kato, S., Fukue, J., & Okazaki, A. T. 1984, *PASJ*, 36, 71
 Molteni, D., Sponholz, H., & Chakrabarti, S. K. 1996, *ApJ*, 457, 805
 Muno, M. P., Morgan, E. H., & Remillard, R. A. 1999, *ApJ*, 527, 321
 Naik, S., Agrawal, P. C., Rao, A. R., Paul, B., et al. 2001, *ApJ*, 546, 1075
 Nakayama, K. 1992, *MNRAS*, 259, 259
 Nakayama, K., & Fukue, J. 1989, *PASJ*, 41, 271
 Nowak, A. M., & Wagoner, R. V. 1991, *ApJ*, 378, 656
 Paczyński, B., & Wiita, P. J. 1980, *A & A*, 88, 23
 Rutledge R E et al. 1999 *ApJ Suppl. Ser.* 124 265
 Sarkar, A., & Das, T. K., 2002, *IJMPD*, Vol. 11. Issue 03, pp. 427 - 432
 Shrader, C. R., & Titarchuk, L. G. 1998, *ApJ*, 499, L31
 Smith, D. M., Heindl, W. A., & Swank, J. H. 2002, *ApJ*, 1, 362
 Titarchuk, L., Lapidus, I., & Muslimov, A. 1998, *ApJ*, 499, 315
 van der Kluis, M. 1989, *ARA & A*, 27, 517
 Yang, R., & Kafatos, M. 1995, *A & A*, 295, 238 (YK)

Figure Captions

Fig. 1: Parameter space division for multi-transonic accretion in four different pseudo-Schwarzschild potentials marked in the figure. Specific angular momentum of the flow (λ) is plotted along the X axis while the flow temperature is plotted along the Y axis in the unit of $10^{10}K$. Note that there is no common set of $\{\lambda, T\}$ for which multi-transonic flow occurs for all potentials, see text for details.

Fig. 2: Multi-transonic isothermal wind as a function of $\{\lambda, T \rightarrow T/T_{10} (= 10^{10}K)\}$ for various black hole potentials marked on the figure. Note how, unlike accretion, multi-transonicity is obtained in wind for very strongly rotating flows (flow with large value of λ).

Fig. 3: Solution topology of multi-transonic shocked accretion along with its ‘self-wind’ for four different black hole potentials marked in the figure. Among the two formal shock solution (marked by vertical lines with downward arrow having *different* values of shock strength, solution through C_1 comes out to be the stable one, see text for detail.

Fig. 4: Global parameter space for accretion solution with shocks (region marked by S_1 and bounded by $B_1C_1D_1$ for the potential Φ_1 and so on) embedded in region for general multi-transonic accretion solutions (region bounded by $A_1B_1C_1$ for Φ_1 and so on) for four different potentials. This figure demonstrates *all* possible $\{\lambda, T \rightarrow T/T_{10} (= 10^{10}K)\}$ for which shock may form in isothermal black hole accretion disc.

Fig. 5: Variation of shock location r_{sh} (scaled in the unit of Schwarzschild radius and plotted along the Z axis) with specific angular momentum (plotted along the X axis) and flow temperature T (scaled in the unit of $T_{10} = 10^{10}K$ and plotted along the Y axis) for four potentials $\Phi_1(a)$, $\Phi_2(b)$, $\Phi_3(c)$ and $\Phi_4(d)$. See text for detail. This figure is complementary to Fig. 4 in the sense that shock locations in this figure are obtained for *all* $\{\lambda, T \rightarrow T/T_{10} (= 10^{10}K)\}$ in Fig. 4 for which shock may form.

Fig. 6: Variation of the amount of energy dissipation $\Delta\epsilon$ at the shock location (plotted along the Z axis) as a function λ (X axis) and $T/T_{10} (= 10^{10}K)$ Y axis for four different potentials $\Phi_1(A)$, $\Phi_2(B)$, $\Phi_3(C)$ and $\Phi_4(D)$, see text for detail.











